This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.



- Hybrid maize seed designated 34N16, representative seed of said hybrid 34N16 1. having been deposited under ATCC accession number
- 2. A maize plant, or its parts, produced by the seed of claim 1.
- 3. Pollen of the plant of claim 2.
- An ovule of the plant of claim 2. 4.
- 5. A tissue culture of regenerable cells of a hybrid maize plant 34N16, representative seed of said hybrid maize plant 34N16 having been deposited under ATCC accession number ____, wherein the tissue regenerates plants capable of expressing all the morphological and physiological characteristics of said hybrid maize plant 34N16.

A tissue culture according to claim 5, the cells or protoplasts being from a tissue selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, silks, flowers, kernels, ears, cobs, husks, and stalks.

7. A maize plant, or its parts, regenerated from the tissue culture of claim 5 and capable of expressing all the morphological and physiological characteristics of hybrid maize plant 34N16, representative seed having been deposited under ATCC accession number

The maize plant of claim 2 wherein said plant is male sterile.

- A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include/employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 2 as a source of said breeding material.
- 10. The maize plant breeding program of claim 9 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing,

11.

15.

pedigree breeding, restriction fragment length polymorphism enhanced selection. genetic marker enhanced selection, and transformation.

A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 2, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay**∖**green, very good drought tolerance, above average early growth, above average test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative maturity of approximately 109 based on the Comparative Relative Maturity Rating System for Narvest moisture of grain.

A hybrid maize plant according to claim 2, wherein the genetic material of said plant contains one or more transgenes.

- 13. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 12 as a source of said breeding material.
- The maize plant breeding program of claim 13 wherein plant breeding techniques 14. are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
 - A maize plant, ohits parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 12, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay green, very good drought tolerance, above average early growth, above average test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative maturity of approximately 109 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

16. A hybrid maize plant according to claim 2, wherein the genetic material of said plant contains one or more genes transferred by backcrossing.

- 17. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising obtaining the maize plant, or its parts, of claim 16 as a source of said breeding material.
- 18. The maize plant breeding program of claim 17 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
- 19. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 16, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay green, very good drought tolerance, above average early growth, above average test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative maturity of approximately 109 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.
- 20. A maize plant, or its parts, having all the morphological and physiological characteristics of the plant of claim 2.

The maize plant of claim 20 wherein said maize plant is male sterile.

- 22. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material comprising: obtaining the maize plant, or its parts, of claim 20 as a source of said breeding material.
- 23. The maize plant breeding program of claim 22 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing,

pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.

24. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 20, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay green, very good drought tolerance, above average early growth, above average test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative matur**i**ty of approximately 109 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

A hybrid maize plant according to claim 20, wherein the genetic material of said plant contains one or more trapsgenes.

A method for developing a maize plant in a maize plant breeding program using 26. plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 25 as a source of said breeding material.

27. The make plant breeding program of claim 26 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.

A maize plant, on tits parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 25, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay green, very good drought tolerance, above average early growth, above average∖test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative maturity of approximately 109 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

28.

32.

29.

A hybrid maize plant according to claim 20, wherein the genetic material of said plant contains one or more genes transferred by backcrossing.

- 30. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: partaining the maize plant, or its parts, of claim 29 as a source of said breeding material.
- 31. The maize plant breeding program of claim 30 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.

A maize plant or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 29, said maize plant capable of expressing a combination of at least two 34N16 traits selected from the group consisting of: high yield potential, with stable yields across yield levels, above average root strength, very good stay green, very good drought tolerance, above average early growth, above average test weight, very good dry down, stronger performance in the Eastern United States, suited to Central corn Belt regions of the United States, and a relative maturity of approximately 109 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

add >